



Ion production rates and cross-sections from the atmospheric observations and comparison with the cloud experiment results

Vladimir Makhmutov, Yuri Stozhkov, Alessandro Franchin, Jasper Kirkby, and CLOUD Collaboration

Citation: [AIP Conference Proceedings](#) **1527**, 342 (2013); doi: 10.1063/1.4803273

View online: <http://dx.doi.org/10.1063/1.4803273>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1527?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Neutrino cross-sections: Experiments](#)

AIP Conf. Proc. **1666**, 060003 (2015); 10.1063/1.4915559

[New anti-neutrino cross-section results from MiniBooNE](#)

AIP Conf. Proc. **1663**, 080003 (2015); 10.1063/1.4919491

[Ionization & Excitation Collision Rates and Cross-Sections in Multicharged Ion Plasmas](#)

AIP Conf. Proc. **808**, 323 (2006); 10.1063/1.2159381

[Josephson cross-sectional model experiment](#)

J. Appl. Phys. **57**, 2550 (1985); 10.1063/1.335444

[On the Evaluation of Rate Coefficients from Limited Cross-Section Data](#)

J. Chem. Phys. **47**, 1894 (1967); 10.1063/1.1712214

Ion Production Rates and Cross-sections from the Atmospheric Observations and Comparison with the Cloud Experiment Results

Vladimir Makhmutov^a, Yuri Stozhkov^a, Alessandro Franchin^b, Jasper Kirkby^c and the CLOUD collaboration

^a *Lebedev Physical Institute RAS, 119991 Moscow, Leninsky prosp. 53, Russia*

^b *University of Helsinki, Department of Physics, FI-00014 Helsinki, Finland*

^c *CERN, CH-1211 Geneva, Switzerland*

Abstract. We present and discuss experimental results obtained from the measurements of cosmic ray fluxes and ion concentrations at different altitudes (from ground level up to 30-35 km) and latitudes (from equator to polar regions) in the Earth's atmosphere. We calculated ion-production cross-sections and ion production rates from these data sets. The same characteristics are possible to be derived from the CLOUD experimental data using ion concentrations, particle beam intensities, etc. We discuss the methods of estimation of these characteristics in the CLOUD experiment.

Keywords: cosmic rays, ionization, cross-section, ion production rate

PACS: 94.20.Qq, 94.30.Hn

INTRODUCTION

In this paper we present short description of the long-term cosmic ray measurements in the atmosphere provided by the Lebedev Physical Institute of the Russian Academy of Sciences. Also, we used experimental data on the light ion measurements in the atmosphere obtained by the Central Aerological Observatory of the Roscomhydromet of the Russia. From these data sets we calculated the ion-production cross-sections (σ) and ion production rates (q) at different altitudes and latitudes in the atmosphere.

The same values σ and q can be defined in the CLOUD experiment where ions are produced by the charged particle fluxes from the accelerator. Finally, we present the results of the comparison ion-production cross-sections and ion production rates obtained in both experiments.

COSMIC RAYS AND IONS MEASUREMENTS IN THE EARTH'S ATMOSPHERE

The galactic cosmic ray (GCR) fluxes are permanent and main natural source of ionizing charge particles in the Earth's atmosphere. The GCR mostly consist of protons (~85-90 % of total flux) and charged nuclei in wide energy range (the 10^6 - 10^{20} eV). When the GCRs propagate through the atmosphere they create secondaries, so called cascade charge particles in wide energy range. These secondary charge

particles are measured regularly in the Earth's atmosphere since 1957 in the long-term cosmic ray experiment carried out by the Lebedev Physical Institute of the Russian Academy of Sciences (LPI RAS) at several locations [1, 2]. As an example, the GCR fluxes time variations at the maximum absorption curve (Pfofzer's maximum) at different latitudes in the Earth's atmosphere are shown in Figure 1.




FIGURE 1. Monthly averages of cosmic ray fluxes in the maximum of absorption curve (Pfofzer's maximum) at different latitudes in the Earth's atmosphere. Curves correspond to the polar northern region (green curve; Murmansk region, geomagnetic cutoff rigidity $R_c=0.5$ GV), southern polar one (blue curve; Mirny station in Antarctica, $R_c=0.04$ GV), middle northern latitude (red curve; Moscow region, $R_c=2.4$ GV) and northern low latitude (light brown; Alma-Ata region, $R_c=6.7$ GV).

We note, that in this experiment the cosmic ray fluxes are recorded at ~ 50 different altitudes in the atmosphere from the ground level up to 30-35 km. Time variation profiles at these altitudes are almost the same, but their amplitudes are decreased when the atmospheric depth is increased.

Ionization of air and ion concentration measurements in the atmosphere were provided in the past in various experiments [3,4]. Figure 2 shows the altitude dependences of light ion concentrations and cosmic ray fluxes as a function of the atmospheric depth. These dependences were obtained at several latitudes with different geomagnetic cutoff rigidities R_c . We note similarity of the curves presented at left and right panels of the Figure 2.




FIGURE 2. The altitude profiles of light ion concentrations (left) and cosmic ray fluxes (right). These dependences were obtained at several locations with different geomagnetic cutoff rigidities R_c (upper numbers in GV).

Monthly averages of cosmic ray fluxes in the maximum of absorption curve (Pfotzer's maximum) at different latitudes in the Earth's atmosphere. These curves correspond to the polar northern region (green curve; Murmansk region, geomagnetic cutoff rigidity $R_c=0.5$ GV), southern polar one (blue curve; Mirny station in Antarctica, $R_c=0.04$ GV), middle northern latitude (red curve; Moscow region, $R_c=2.4$ GV) and northern low latitude (light brown; Alma-Ata region, $R_c=6.7$ GV).

Comparative analysis of results of the CRs and ions measurements in the atmosphere allowed us to calculate the ion-production cross-sections and ion production rates at different altitudes.

CLOUD EXPERIMENT: CHARGE PARTICLE FLUXES AND IONS

In the CLOUD experiment charged particles (3.5 GeV/c pions) from CERN accelerator PS215 penetrate into the CLOUD chamber (volume $\sim 26\text{m}^3$) and generate ions inside the chamber [5]. The scintillation hodoscope provide measurements of the space and time distributions of charged particle flux. The experimental devices NAIS and APITOFF provide continuous records of the positive and negative ion concentrations in the CLOUD chamber. We used these data sets to calculate the ion-production cross-sections and ion production rates in the chamber under given conditions. We discuss method of the σ and q value calculation and compare results obtained in the atmosphere and in the CLOUD experiment.

ACKNOWLEDGMENTS

We would like to thank CERN for supporting CLOUD with important technical and financial resources, and for providing a particle beam from the CERN Proton Synchrotron. This research has received funding from the EC Seventh Framework Programme (Marie Curie Initial Training Network "CLOUD-ITN" no. 215072, MC-ITN "CLOUD-TRAIN" no. 316662, and ERC-Advanced "ATMNUCLE" grant no. 227463), the German Federal Ministry of Education and Research (project nos. 01LK0902A and 01LK1222A), the Swiss National Science Foundation (project nos. 20

0020_135307 and 206620_130527), the Academy of Finland (Center of Excellence project no. 1118615), the Academy of Finland (135054, 133872, 251427, 139656, 139995, 137749, 141217, 141451), the Finnish Funding Agency for Technology and Innovation, the Nessling Foundation, the Austrian Science Fund (FWF; project no. P19546 and L593), the Portuguese Foundation for Science and Technology (project no. CERN/FP/116387/2010), the Swedish Research Council, Vetenskapsrådet (grant 2011-5120), the Presidium of the Russian Academy of Sciences and Russian Foundation for Basic Research (grant 12-02-91522-CERN), and the U.S. National Science Foundation (grants AGS1136479 and CHE1012293).

REFERENCES

1. A. N. Charakhchyan: Investigation of stratosphere cosmic ray intensity fluctuations induced by processes on the Sun, *Soviet Physics Uspekhi* **7**, 358–374 (1964).
2. Y. I. Stozhkov, N. S. Svirzhevsky, G. A. Bazilevskaya, M. B. Krainev, A. K. Svirzhevskaya, V. S. Makhmutov, V. I. Logachev, and E. V. Vashenyuk, *Astrophys. Space Sci. Trans.* **7**, 379–382 (2011).
3. H.V. Neher, Cosmic rays at high latitudes and altitudes covering four solar maxima, *J. Geophys. Res.*, **76**, 1637-1651 (1971).
4. V.I. Ermakov, G.A. Bazilevskaya, P.E. Pokrevsy, Y.I. Stozhkov, Ion balance equation in the atmosphere, *J. Geophys. Res.*, **102**, 23413-23419 (1997).
5. J. Kirkby, J. Curtius, J. Almeida, et al., Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, **476**, doi:10.1038 (2011).